

Poster presentation

## Design of novel semisynthetic metalloenzyme from thermolysin

Mohd Basyaruddin Abdul Rahman\*<sup>1</sup>, Ahmad Haniff Jaafar<sup>1</sup>, Mahiran Basri<sup>1</sup>, Raja Noor Zaliha Raja Abdul Rahman<sup>2</sup>, Abu Bakar Salleh<sup>2</sup> and Habibah Abdul Wahab<sup>3</sup>

Address: <sup>1</sup>Department of Chemistry, Faculty of Science, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia, <sup>2</sup>Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia and <sup>3</sup>Laboratory of Biocrystallography and Bioinformatic Structure, Universiti Sains Malaysia, 11800 Minden, Penang, Malaysia

Email: Mohd Basyaruddin Abdul Rahman\* - basya@science.upm.edu.my

\* Corresponding author

from BioSysBio 2007: Systems Biology, Bioinformatics and Synthetic Biology  
Manchester, UK. 11–13 January 2007

Published: 8 May 2007

BMC Systems Biology 2007, 1(Suppl 1):P68 doi:10.1186/1752-0509-1-S1-P68

This abstract is available from: <http://www.biomedcentral.com/1752-0509/1?issue=S1>

© 2007 Rahman et al; licensee BioMed Central Ltd.

### Background

Initial applications of biocatalysis involved the used of naturally occurring enzyme. With new challenges in green chemical reaction, biocatalyst that shed the light is metalloenzyme, which function as enzyme and contain metal that are tightly attached and always isolated with the protein [1]. In recent years, enzyme engineering has proven to be an invaluable tool for elucidating biocatalytic mechanisms as well as producing enzymes for industrial purposes. Approaches developed for *in vivo* chemical modification and *in silico* computational methods promise to increase the scope and have already been used successfully to alter existing protein so that they have better stability and functionality [2]. This task might be good to address in designing a new biocatalyst with improved properties.

### Methods

The AutoDock programme 3.05 was employed in order to identify the binding conformations of the ligands and the metal ions and to perform docking using Lamarckian Genetic Algorithm (LGA) [3]. The coordinate of thermolysin-substrate free structure coded as 1KEI was taken from Brookhaven Protein Data Bank (PDB).

### Results

The predicted KEI-ligand complexes with the lowest final docked energy for PSE and PHN were -6.71 kcal/mol at pocket 45 and -6.60 kcal/mol at pocket 47, respectively. Non-covalent interactions of hydrogen bond and hydrophobic interaction between protein and ligands established the final conformation. Analysis on finding the most favorable metal ions to dock onto each complex found that Mg<sup>2+</sup> was docked onto KEI-PSE45 complex with final docked energy of -1.09 kcal/mol and performed four interactions with the PSE ligand. Meanwhile, Ca<sup>2+</sup> represented the best metal ions to dock to the KEI-PHN47 complex with final docked energy of -4.12 kcal/mol and performed three interactions with the nearby residues.

### Conclusion

An important branch of novel protein design is through engineering and design of new metal-binding sites into native proteins. By employing *in silico* approach of molecular docking, screening of putative ligand for possible interactions may enhance the discovery of novel semisynthetic enzyme and lead to a new protein function. Finally, the framework which was introduced for the experiment may be a competent method for screening potential metal ions in this *in vivo* route.

## References

1. Davies RR, Kuang H, Qi D, Mazhary A, Mayaan E, Distefeno MD: **Artificial Metalloenzymes Based on Protein Cavities: Exploring the Effect of Altering the Metal Ligand Attachment Position by Site Directed Mutagenesis.** *Bioorg Med Chem Lett* 1999, **9**:79-84.
2. Abdul Rahman MB, Misran A, Abdul Wahab H, Abdul Rahman RNZ, Salleh AB, Basri M: **Screening and Docking of Chemical Ligands onto Pocket Cavities of Protease for Designing a Biocatalyst.** *Biocatal Biotransform* 2005, **23**:211-216.
3. Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK, Olson AJ: **Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function.** *J Comput Chem* 1998, **19**:1639-1662.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

